

Evaluation of the low-lying energy levels of two- and three-electron configurations for multi-charged ions

O. Yu. Andreev*,¹ L. N. Labzowsky,¹ G. Plunien,² and G. Soff²

¹ *Institute of Physics, St. Petersburg State University, Ulyanovskaya ul. 1, 198904, Petershof, St. Petersburg, Russia*

² *Institut für Theoretische Physik, Technische Universität Dresden, Mommsenstraße 13, D-01062, Dresden, Germany*
(February 2, 2008)

Accurate QED evaluations of the one- and two-photon interelectron interaction for low lying two- and three-electron configurations for ions with nuclear charge numbers $60 \leq Z \leq 93$ are performed. The three-photon interaction is also partly taken into account. The Coulomb gauge is employed. The results are compared with available experimental data and with different calculations. A detailed investigation of the behaviour of the energy levels of the configurations $1s_{1/2}2s_{1/2}^1S_0$, $1s_{1/2}2p_{1/2}^3P_0$ near the crossing points $Z = 64$ and $Z = 92$ is carried out. The crossing points are important for the future experimental search for parity nonconserving (PNC) effects in highly charged ions.

PACS number(s): 31.30.Jv, 31.10.+z

I. INTRODUCTION

During the most recent years the energy levels of two- and three-electron configurations are under very intensive experimental and theoretical investigation.

Accurate calculations of the energy levels for the two-electron configurations $1s_{1/2}2s_{1/2}^1S_0$, $1s_{1/2}2p_{1/2}^3P_0$, and $1s_{1/2}2s_{1/2}^3S_1$ were already performed in [1,2]. In these papers the interelectron interaction has been considered in various approximations: on the basis of variational Schrödinger-wave functions with inclusion of relativistic correlations [1] and using relativistic many-body perturbation theory (RMBPT) [2]. For a long time these approaches defined the standard concerning the level of accuracy. In recent years due to new developments in experimental and theoretical methods the necessity to improve the accuracy of calculations became urgent. Recently, rigorous QED evaluations of two-photon exchange corrections for low lying configurations [3,4] have been performed.

First calculations of the energy levels for the three-electron configuration have been presented in [5–9]. As in the early papers on two-electron ions in these calculations the two- and many-photon exchange has been considered approximately. Complete QED calculations of the two-photon exchange for three-electron ions have been performed in [10,4]. Within the framework of RMBPT the three-photon exchange has also been taken into account in [11,10,4].

In this paper we present an extension of previous calculations for two- and three-electron ions [4] for a variety of Z values. Here we include also the evaluations for the $1s_{1/2}2p_{1/2}^3P_0$ level, which have not been performed in [4]. Special attention has been paid to elaborate the behaviour of the $1s_{1/2}2s_{1/2}^1S_0$ and $1s_{1/2}2p_{1/2}^3P_0$ levels near $Z = 64$ and $Z = 92$, where they become very close to each other. Accordingly, these system become suitable for monitoring parity-nonconserving (PNC) effects. Experimental investigations of PNC effects in two-electron highly charged ions are under intensive discussion [12–15]. The preparation of such experiments requires a precise knowledge of the exact difference between these energy levels.

II. THEORY

In this article we evaluate corrections to the energy levels due to photon exchange. To calculate these corrections we employ the adiabatic S -matrix approach [16,17] and the line profile approach (LPA) [18]. Both methods are based on the Furry picture [19], which describes the many-electron atom as a set of bound electrons, moving in the external field of the nucleus and interacting with each other via the exchange of photons. With the aid of the Feynman rules for bound-state QED [20,21] the processes giving rise to corrections to the energy levels can be represented in terms of Feynman graphs. The photon-exchange corrections evaluated in this article are depicted in Figs. 1-5.

As it has been shown in Refs. [22,20] the S -matrix approach is best suited for the evaluation of corrections described by the irreducible parts of the diagrams. However, its application to the evaluation of the reducible parts becomes

*Email: oleg@landau.phys.spbu.ru

rather complicated. Accordingly, for calculating the reducible parts we utilized the LPA. For a detailed description and for the application of this method we refer to [4].

The Coulomb gauge will be used throughout this paper. The photon propagators for Coulomb ($g = c$) and transverse ($g = t$) photons are given by [20,21]:

$$D_{\mu_1\mu_2}^g(x_1, x_2) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} d\Omega I_{\mu_1\mu_2}^g(\Omega, r_{12}) e^{i\Omega(t_1-t_2)} \quad (1)$$

together with the temporal Fourier transforms

$$I_{\mu_1\mu_2}^c(\Omega, r_{12}) = \frac{\delta_{\mu_1 4} \delta_{\mu_2 4}}{r_{12}} \quad (2)$$

and

$$I_{\mu_1\mu_2}^t(\Omega, r_{12}) = \left(\frac{\delta_{\mu_1\mu_2}}{r_{12}} e^{i|\Omega|r_{12}} + \nabla_{1\mu_1} \nabla_{2\mu_2} \frac{1}{r_{12}} \frac{1 - e^{i|\Omega|r_{12}}}{|\Omega|^2} \right) (1 - \delta_{\mu_1 4})(1 - \delta_{\mu_2 4}). \quad (3)$$

A. The two-electron configurations

At first we consider the photon exchange corrections for two-electron configurations. The wavefunction of a two-electron configurations is represented by

$$\Psi_{JMj_1j_2l_1l_2}(\mathbf{r}_1, \mathbf{r}_2) = N \sum_{m_1m_2} C_{JM}^{j_1j_2}(m_1m_2) [\psi_{j_1l_1m_1}(\mathbf{r}_1)\psi_{j_2l_2m_2}(\mathbf{r}_2) - \psi_{j_1l_1m_1}(\mathbf{r}_2)\psi_{j_2l_2m_2}(\mathbf{r}_1)], \quad (4)$$

where $N = 1/2$ for equivalent electrons and $N = 1/\sqrt{2}$ for non-equivalent electrons, $C_{JM}^{j_1j_2}(m_1m_2)$ is a Clebsch-Gordan symbol. By means of Eq. (4) we can specify the configuration $1s2s^3S_1$ by setting $a, b = 1s_+, 2s_+$, where $\pm$ denote the two different projections of the total electron angular momentum and we can derive the energy corrections according to the formula:

$$\Delta E(1s2s^3S_1) = F_{1s_+2s_+;1s_+2s_+}, \quad (5)$$

$$F_{ab;cd} = F_{abcd} - F_{bacd}. \quad (6)$$

Here $F_{ab\dots}$ denotes a function of one-electron states which are described by wave functions $\psi_a, \psi_b, \dots$. The form of the function F depends on the type of Feynman graph under consideration (see below). For the electron configurations $1s2s^1S_0$ and $1s2p^1P_0$ the energy corrections are given by

$$\Delta E(1s2s^1S_0) = F_{1s_-2s_+;1s_-2s_+} - F_{1s_+2s_-;1s_-2s_+}, \quad (7)$$

and

$$\Delta E(1s2p^3P_0) = F_{1s_-2p_+;1s_-2p_+} - F_{1s_+2p_-;1s_-2p_+}, \quad (8)$$

respectively. The corrections due to one-photon exchange are represented by the graph in Fig.1. This diagram is irreducible so that the S -matrix approach can be applied leading to

$$F_{a'b'ab}^{(1)} = \sum_g I^g(\varepsilon_{a'} - \varepsilon_a)_{a'b'ab}. \quad (9)$$

Here we have introduced the following notation (see definitions Eqs. (2) and (3))

$$I_{a'b'ab}^g(\Omega) \equiv \sum_{\mu_1\mu_2} \int \bar{\psi}_{a'}(\mathbf{r}_1) \bar{\psi}_{b'}(\mathbf{r}_2) \gamma_{\mu_1}^{(1)} \gamma_{\mu_2}^{(2)} I_{\mu_1\mu_2}^g(\Omega, r_{12}) \psi_a(\mathbf{r}_1) \psi_b(\mathbf{r}_2) d^3r_1 d^3r_2, \quad (10)$$

where the Dirac matrices $\gamma_{\mu_i}^{(i)}$ are acting on wave functions depending on spatial variables $\mathbf{r}_i$, respectively. For $g = c$ Eq. (9) determines the first-order Coulomb correction while for $g = t$ we obtain the first-order Breit correction.

The two-photon exchange corrections are represented by the graphs in Fig. 2. The “box” diagram is reducible. Its reducible part is defined by the condition $\varepsilon_{n_1} + \varepsilon_{n_2} = \varepsilon_a + \varepsilon_b$. The “cross” diagram is irreducible. However, it is most convenient to extract the contribution with n_1, n_2 equal to a or b and to treat it like a reducible part of the “cross” diagram. Contributions due to states n_1, n_2 included in the reducible parts are called reference state contribution. Application of the S -matrix approach for calculating the irreducible part and of the LPA for the reducible part of the “box” and “cross” diagrams, respectively, results in the explicit formulas:

$$F_{a'b'ab}^{(2)(\text{box},\text{irr})} = \sum_{\text{gg}'} \sum'_{n_1 n_2} \left\{ \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{a'b'n_1 n_2} I^{g'}(\Omega - \varepsilon_{a'} + \varepsilon_a)_{n_1 n_2 ab}}{(\varepsilon_a + \varepsilon_b - \varepsilon_{n_1} - \varepsilon_{n_2})(\Omega - \varepsilon_{n_2} + \varepsilon_{b'} + i0 \varepsilon_{n_2})} d\Omega \right. \\ \left. + \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{b'a'n_1 n_2} I^{g'}(\Omega - \varepsilon_a + \varepsilon_{a'})_{n_1 n_2 ba}}{(\varepsilon_a + \varepsilon_b - \varepsilon_{n_1} - \varepsilon_{n_2})(\Omega - \varepsilon_{n_2} + \varepsilon_{a'} + i0 \varepsilon_{n_2})} d\Omega \right\}, \quad (11)$$

$$F_{a'b'ab}^{(2)(\text{box},\text{red})} = -\frac{1}{2} \sum_{\text{gg}'} \sum''_{n_1 n_2} \left\{ \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{a'b'n_1 n_2} I^{g'}(\Omega - \varepsilon_{a'} + \varepsilon_a)_{n_1 n_2 ab}}{(\Omega - \varepsilon_{n_2} + \varepsilon_{b'} + i0 \varepsilon_{n_2})^2} d\Omega \right. \\ \left. + \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{b'a'n_1 n_2} I^{g'}(\Omega - \varepsilon_a + \varepsilon_{a'})_{n_1 n_2 ba}}{(\Omega - \varepsilon_{n_2} + \varepsilon_{a'} + i0 \varepsilon_{n_2})^2} d\Omega \right\}, \quad (12)$$

$$F_{a'b'ab}^{(2)(\text{cross},\text{irr})} = \sum_{\text{gg}'} \sum'_{n_1 n_2} \left\{ \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{b'n_2 n_1 a} I^{g'}(\Omega - \varepsilon_{a'} + \varepsilon_a)_{n_1 a' b n_2}}{(\varepsilon_{n_2} - \varepsilon_{n_1} - \varepsilon_a + \varepsilon_{b'}) (\Omega - \varepsilon_{n_2} + \varepsilon_a + i0 \varepsilon_{n_2})} d\Omega \right. \\ \left. + \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{n_1 b' a n_2} I^{g'}(\Omega - \varepsilon_a + \varepsilon_{a'})_{a' n_2 n_1 b}}{(\varepsilon_{n_2} - \varepsilon_{n_1} + \varepsilon_a - \varepsilon_{b'}) (\Omega - \varepsilon_{n_2} + \varepsilon_{b'} + i0 \varepsilon_{n_2})} d\Omega \right\}, \quad (13)$$

$$F_{a'b'ab}^{(2)(\text{cross},\text{red})} = \sum_{\text{gg}'} \sum''_{n_1 n_2} \frac{i}{2\pi} \int_{-\infty}^{\infty} \frac{I^g(\Omega)_{b'n_2 n_1 a} I^{g'}(\Omega - \varepsilon_{a'} + \varepsilon_a)_{n_1 a' b n_2}}{(\Omega - \varepsilon_{n_2} + \varepsilon_a + i0 \varepsilon_{n_2})^2} d\Omega. \quad (14)$$

The prime at the summation symbols indicates that the proper reference state members are ejected. The double prime indicates that only the reference state members are retained. In order to avoid division by zero in Eq. (13) in the case $a = b'$ and $n_1 = n_2$ one has to take the limit $\varepsilon_{n_1} \rightarrow \varepsilon_{n_2}$ in both terms on the right-hand side. Thus the singularities cancel. It should be stressed that Eq. (14) coincides with the result of this limiting process. From Eqs. (12) and (14) it follows automatically, that the corrections vanish for $g = g' = c$. The case $g = g' = c$ corresponds to the Coulomb-Coulomb correction, the case $g = g' = t$ determines the Breit-Breit and the case $g = c, g' = t$ or $g = t, g' = c$ refers to the Coulomb-Breit interaction.

For high- Z ions considered in this work the third-order contribution turns out to be small and it is sufficient to take into account its dominant part only. Accordingly, we consider only the third-order Coulomb and unretarded Breit “box” corrections. The corresponding Feynman graph is displayed in Fig. 3. The formula for the irreducible part of the third-order “box” correction can be derived in the same manner as for the corrections given by Eqs. (9, 11). It takes the form

$$F_{a'b'ab}^{(3)(\text{box},\text{irr})} = \sum_{\text{gg}'} \sum'_{g'' n_1 n_2 n_3 n_4} \frac{I^g_{a'b'n_3 n_4} I^{g'}_{n_3 n_4 n_1 n_2} I^{g''}_{n_1 n_2 ab}}{(\varepsilon_{n_3} + \varepsilon_{n_4} - \varepsilon_{a'} - \varepsilon_{b'}) (\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b)}, \quad (15)$$

where the prime indicates that the reference state contributions are excluded from the summation. Here the reference states are defined by the conditions $\varepsilon_{n_1} + \varepsilon_{n_2} = \varepsilon_a + \varepsilon_b$ or $\varepsilon_{n_3} + \varepsilon_{n_4} = \varepsilon_a + \varepsilon_b$. Applying the LPA to the graph in Fig. 3 we derive the following expression for the reducible part

$$F_{a'b'ab}^{(3)(\text{box},\text{red})} = \sum_{\text{gg}'} \sum''_{g'' n_1 n_2 n_3 n_4} I^g_{a'b'n_3 n_4} I^{g'}_{n_3 n_4 n_1 n_2} I^{g''}_{n_1 n_2 ab} \\ \times \left\{ \frac{(-1)}{2(\varepsilon_{n_3} + \varepsilon_{n_4} - \varepsilon_{a'} - \varepsilon_{b'})^2} + \frac{(-1)}{2(\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b)^2} \right\}, \quad (16)$$

where the double prime indicates that the summation is running only over the reference states. The terms leading to vanishing denominators in Eq. (16) should be omitted.

B. The three-electron configurations

Now we turn to three-electron ions. Here we consider three-electron configurations with a closed $(1s)^2$ shell, which can be described by the wave function

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = \frac{1}{\sqrt{3!}} \sum_{i,j,k=1,2,3} \epsilon_{ijk} \psi_i(\mathbf{r}_1) \psi_j(\mathbf{r}_2) \psi_k(\mathbf{r}_3). \quad (17)$$

ϵ_{ijk} denotes the Levi-Civita symbol and $\psi_1(\mathbf{r})$, $\psi_2(\mathbf{r})$, $\psi_3(\mathbf{r})$ denote one-electron wave functions.

As in the two-electron case we have to consider corrections represented by the two-electron Feynman graphs depicted in Figs. 1-3. Their contribution to the energy shift is given by

$$\Delta E(\{abc\}) = F_{ab;ab} + F_{bc;bc} + F_{ca;ca}, \quad (18)$$

where $F_{ab;cd}$ is given by Eqs. (10) and (9-16). The set $\{abc\}$ is equal to the set $\{1s_+, 1s_-, 2s_{1/2+}\}$ for the configuration $(1s)^2 2s_{1/2}$ and to the set $\{1s_+, 1s_-, 2p_{1/2+}\}$ for the configuration $(1s)^2 2p_{1/2}$, respectively. The symbol $\pm$ refers to the different angular momentum projections.

Besides the two-electron diagrams, in three-electron problem we have to take into account the additional three-electron Feynman graphs depicted in Figs. 4 and 5. The contribution of the three-electron graphs can be calculated according to

$$\Delta E(\{abc\}) = \sum_{\substack{i',j',k'=1,2,3 \\ i,j,k=1,2,3}} \epsilon_{i'j'k'} \epsilon_{ijk} F_{i'j'k'ijk}, \quad (19)$$

where the indices 1, 2, 3 at F symbol must be replaced by a, b, c respectively, i.e. $F_{ab;cab} \equiv F_{123123}$, etc. Eq. (19) includes the contribution of the “direct” and all possible “exchange” diagrams which occur in the three-electron case.

Expressions for $F_{a'b'c'abc}$ corresponding to the graph in Fig. 4 are

$$F_{a'b'c'abc}^{(2)(\text{step,irr})} = \sum_{gg'} \sum_n' \frac{I^g(\varepsilon_a - \varepsilon_{a'})_{na'ba} I^{g'}(\varepsilon_{c'} - \varepsilon_c)_{b'c'nc}}{\varepsilon_a + \varepsilon_b - \varepsilon_{a'} - \varepsilon_n}, \quad (20)$$

$$F_{a'b'c'abc}^{(2)(\text{step,red})} = \sum_{gg'} \sum_n'' \frac{\partial}{\partial \omega} \left[I^g(\varepsilon_a - \varepsilon_{a'} + \omega)_{na'ba} I^{g'}(\varepsilon_{c'} - \varepsilon_c + \omega)_{b'c'nc} \right] \Big|_{\omega=0}, \quad (21)$$

where the prime at the summation symbol indicates that the summation runs over all n except for the case when the set of one-electron states $\{a', n, c\}$ is equivalent to the set $\{a, b, c\}$. The latter refers to reference states. The double prime implies that the summation runs over the reference states only. As for the two-electron contributions we have here $g, g' = c, t$. No reducible contribution arises for $g = g' = c$.

As it has been mentioned above, for the three-photon corrections we take into account only their dominant parts, i.e., the third-order Coulomb and unretarded Breit “box” contributions. The corresponding tree-electron Feynman graphs are displayed in Fig. 5. The formulas for the irreducible and the reducible parts of the third-order “box” correction are derived in the same manner as in Eqs. (15,16). The irreducible part can be expressed as

$$\begin{aligned} F_{a'b'c'abc}^{(3)(\text{step-box,irr})} = & \sum_{gg'g''} \sum_{n_1 n_2 n_3}' \frac{I_{a'b'n_1 n_3}^g I_{n_3 c' n_2 c}^{g'} I_{n_1 n_2 ab}^{g''}}{(\varepsilon_{n_1} + \varepsilon_{n_3} - \varepsilon_{a'} - \varepsilon_{b'}) (\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b)} \\ & + 2 \sum_{gg'g''} \sum_{n_1 n_2 n_3}' \frac{I_{b'c'n_3 c}^g I_{a'n_3 n_1 n_2}^{g'} I_{n_1 n_2 ab}^{g''}}{(\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b) (\varepsilon_{n_3} + \varepsilon_{a'} - \varepsilon_a - \varepsilon_b)} \\ & + \sum_{gg'g''} \sum_{n_1 n_2 n_3}' \frac{I_{a'c'n_1 n_3}^g I_{b'n_3 n_2 c}^{g'} I_{n_1 n_2 ab}^{g''}}{(\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b) (\varepsilon_{n_1} + \varepsilon_{n_3} - \varepsilon_{a'} - \varepsilon_{c'})}, \end{aligned} \quad (22)$$

where the prime at the summation symbols indicates that the first summation does not run over states for which either the set $\{n_1, n_2, c\}$ or the set $\{n_1, n_3, c'\}$ are equivalent to the set $\{a, b, c\}$; the second summation does not run over the states for which the sets $\{n_1, n_2, c\}$ or $\{a', n_3, c\}$ are equivalent to the set $\{a, b, c\}$ and the third summation does not run over the states for which the sets $\{n_1, n_2, c\}$ or $\{n_1, n_3, b'\}$ are equivalent to the set $\{a, b, c\}$ (the cases of reference states). The reducible part of the third-order “step-box” corrections (see Fig. 5) can be cast into the form

$$\begin{aligned}
F_{a'b'c'a \ b \ c}^{(3)(\text{step-box,red})} = & \sum_{gg'g''} \sum_{n_1 n_2 n_3}'' I_{a'b'n_1 n_3}^g I_{n_3 c' n_2 c}^{g'} I_{n_1 n_2 a b}^{g''} \\
& \times \left\{ \frac{(-1)}{2(\varepsilon_{n_1} + \varepsilon_{n_3} - \varepsilon_{a'} - \varepsilon_{b'})^2} + \frac{(-1)}{2(\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b)^2} \right\} \\
& + 2 \sum_{n_1 n_2 n_3}'' I_{b'c'n_3 c}^g I_{a'n_3 n_1 n_2}^{g'} I_{n_1 n_2 a b}^{g''} \\
& \times \left\{ \frac{(-1)}{2(\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b)^2} + \frac{(-1)}{2(\varepsilon_{n_3} + \varepsilon_{a'} - \varepsilon_a - \varepsilon_b)^2} \right\} \\
& + \sum_{n_1 n_2 n_3}'' I_{a'c'n_1 n_3}^g I_{b'n_3 n_2 c}^{g'} I_{n_1 n_2 a b}^{g''} \\
& \times \left\{ \frac{(-1)}{2(\varepsilon_{n_1} + \varepsilon_{n_2} - \varepsilon_a - \varepsilon_b)^2} + \frac{(-1)}{2(\varepsilon_{n_1} + \varepsilon_{n_3} - \varepsilon_{a'} - \varepsilon_{c'})^2} \right\}, \tag{23}
\end{aligned}$$

where the double prime at the summation symbols indicates that the summations run over the corresponding reference states only (see the explanations for Eq. (22)). It becomes evident that the contributions due to the graphs Fig. 5b, c are equal. Therefore, we account for them by taking twice the contribution of the graph Fig. 5b.

III. NUMERICAL RESULTS AND DISCUSSION

The major result of the present work consists in the calculation of the two- and three-photon exchange corrections to the energy levels of two-electron configurations 2^1S_0 , 2^3P_0 , 2^3S_1 and three-electron configurations $(1s)^2 2s_{1/2}$, $(1s)^2 2p_{1/2}$. The two-photon exchange correction represents the leading part of the perturbation theory in second order. Accordingly, the main uncertainty of the theoretical values calculated earlier has been due to this correction.

In order to represent the Coulomb potential of the nucleus we employ a Fermi model for the nuclear density distribution

$$\rho(r) = \frac{N}{1 + \exp[(r - c)/a]}, \tag{24}$$

where N is a normalization constant, $a = 0.5350$ fm and c is deduced via the equation

$$4\pi \int_0^\infty \rho(r) r^4 dr = \langle r^2 \rangle, \tag{25}$$

where $\langle r^2 \rangle^{1/2}$ is the root-mean-square nuclear radius. In Table I we also display the values for the nuclear root-mean-square radii employed in this work. They have been taken from [23]. For nuclei with charge numbers Z not presented in [23] we utilize the empirical formula [24]

$$\langle r^2 \rangle^{1/2} = (0.836 A^{1/3} + 0.570) \text{ fm}, \tag{26}$$

where A is the atomic mass number.

The results of our calculation of the two-photon exchange correction are presented in Tables II, III, IV, V, VI, VII for two-electron configurations and in Tables VIII, IX, X, XI for three-electron configurations, respectively. Our calculation is performed rigorously within the framework of QED. For reasons of clarity the corresponding corrections referring to contributions of the Feynman graphs in Figs. 2 and 4 are also listed separately in these tables. For details concerning the numerical procedure we refer to [4]. The accuracy of the present calculations is on the level of about 0.0001 a.u.

We also have taken into account the dominant part of the three-photon exchange correction. Details of the approximation made were given in Sec. II. The results of the calculation for the three-photon exchange correction

are presented in Tables XII, XIII, XIV, XV, XVI, XVII for two-electron configurations and in Tables XVIII, XIX, XX, XXI for three-electron configurations, respectively. Again, the contribution due to the exchange of the various photons are compiled separately in the tables. The correction caused by the exchange of three Breit photons is not included since it was found to be less than 0.001 eV. In view of the approximation used to evaluate the three-photon exchange correction these values are given within an inaccuracy of about 10 % [4].

In Tables XXII, XXIII, XXIV, XXV, XXVI, XXVII, XXVIII, XXIX, XXX we collect all available corrections to the energy levels of the two-electron configurations under consideration. In order to compare our numerical data for the two-electron configurations 2^3P_0 and 2^3S_1 with other results in the literature we also provide values for the two-photon exchange correction as it has been derived in [3]. We find that the data presented in [3] deviate from our results by not more than 0.0003 a.u.

Values for the energy of the three-electron configurations are presented in Tables XXXI, XXXII, XXXIII, XXXIV, XXXV, XXXVI. Comparing our results for the two-photon exchange correction with data presented in Ref. [10], we achieved a very good agreement for the $(1s)^22s_{1/2}$ configuration. However, for the $(1s)^22p_{1/2}$ level we find a discrepancy of about 0.0035 a.u. for $Z = 60, 70$. The three-photon exchange correction is compared with the results obtained in [11]. In [11] the exchange of two and three Breit photons has been neglected.

The numbers for the recoil correction included in the tables for the total level energies of two- and three-electron configurations are obtained by interpolation for those Z values not calculated in the referred paper. The data for the self-energy (SE) screening and vacuum polarization (VP) screening of three-electron configurations as well as for the VP screening of two-electron configurations have been obtained via a similar interpolation. Results for the SE screening corrections of two-electron configurations have been obtained according to a procedure which is based on the results provided in [25]. In particular we refer to Table II of Ref. [25], where the self-energy screening function $f(Z\alpha)$ for K- and L-shell single-electron states have been presented. From these values one can deduce the corresponding self-energy shift of a single-electron state due to the screening effect of another single electron state. E.g., we may denote by $E^{1s \text{ by } 2s}$ the screening correction to the $1s$ -electron self-energy shift due to the $2s$ -electron state and by $E^{2s \text{ by } 1s}$ the screening correction to the $2s$ -self energy originating from the $1s$ -electron state, respectively. Accordingly, we suppose that the sum of the SE screening corrections $E^{1S_0 \text{ scr}}$ and $E^{3S_1 \text{ scr}}$ for the $1S_0$ and $3S_1$ configuration is represented by $E^{1s \text{ by } 2s} + E^{2s \text{ by } 1s} = E^{1S_0 \text{ scr}} + E^{3S_1 \text{ scr}}$. Then we suppose that $E^{1S_0 \text{ exch}}/E^{3S_1 \text{ exch}} = E^{1S_0 \text{ scr}}/E^{3S_1 \text{ scr}}$, where $E^{1S_0 \text{ exch}}$ and $E^{3S_1 \text{ exch}}$ are the first-order interelectron interaction corrections for the corresponding configurations. For the $3P_0$ configuration we define $E^{1s \text{ by } 2p_{1/2}} + E^{2p_{1/2} \text{ by } 1s} = E^{3P_0 \text{ scr}}$.

In Table XXXVII we present the total values for the energy levels of the two-electron configurations derived in this paper and compare them with the results of [2] and [1], respectively. The differences between the energy levels are also listed in that table. We should note that in Ref. [1,2], different approaches have been employed, i.e., the relativistic-all-order-theory (AO) [2] and the unified theory (UT) [1]. Compared with the rigorous QED approach, these theories involve several approximations, i.e., neglect of (i) negative energy states, (ii) crossed photon contributions, and (iii) exact retardation effects. However, they account for some part of higher-order interelectron interaction corrections. Accordingly, for highly-charged ions the total data derived in Tables XXII, XXIII, XXV, XXVI, XXVIII, XXIX, XXXI, XXXII, XXXIV, XXXV provide the most accurate theoretical predictions for the energy levels at present.

From the results presented in Table XXXVII one can conclude that the configurations 2^1S_0 and 2^3P_0 cross within the interval $60 < Z < 70$. Experimental investigation of PNC effects in heliumlike ions requires a precise knowledge of the energy difference between these levels at $Z = 63$ [15]. UT theory [1] predicts a value for this difference of about 0.168 eV, while the calculation presented in this paper gives a larger value of 0.593 eV. However, our calculations predict that the crossing of these levels takes place near $Z = 66$ with an energy difference of about -0.016 eV. Nevertheless, the He-like Eu ion ($Z = 63$) seems most suitable for the search of PNC effects [15]. We also investigated the splitting $E(2^1S_0) - E(2^3P_0)$ for two isotopes $^{151}_{63}\text{Eu}$ and $^{153}_{63}\text{Eu}$ and obtained an energy difference 0.001 eV, which does not change the conclusions made in [15]. The present calculation also indicates that the other crossing point can be expected to be close to $Z = 89, 90$.

IV. ACKNOWLEDGEMENTS

The authors are indebted to Prof. W. Nagel from the computer center of the TU Dresden for providing access to the all necessary computer facilities. O.A. is grateful to the TU Dresden for the hospitality during his visits in 2001 and 2002 and to the DFG for financial support. The work of O.A. and L.N. was supported by the RFBR Grant No. 02-02-16578 and by Minobrazovanie grant No. E00-3.1-7. G.P. and G.S. acknowledge financial support from BMBF, DFG and GSI.

- [1] G. W. Drake, Can. J. Phys. **66**, 586 (1988).
- [2] D. R. Plante, W. R. Johnson, and J. Sapirstein, Phys. Rev. A **49**, 3519 (1994).
- [3] P. J. Mohr and J. Sapirstein, Phys. Rev. A **62**, 052501 (2000).
- [4] O. Yu. Andreev, L. N. Labzowsky, G. Plunien, and G. Soff, Phys. Rev. A **64**, 042513 (2001).
- [5] W. R. Johnson, S. A. Blundell, and J. Sapirstein, Phys. Rev. A **37**, 2764 (1988).
- [6] Y. K. Kim, D. H. Baik, P. Indelicato, and J. P. Desclaux, Phys. Rev. A **44**, 148 (1991).
- [7] S. Blundell, P. J. Mohr, W. R. Johnson, and J. Sapirstein, Phys. Rev. A **48**, 2615 (1993).
- [8] M. H. Chen, K. T. Cheng, W. R. Johnson, and J. Sapirstein, Phys. Rev. A **52**, 266 (1995).
- [9] H. Persson, I. Lindgren, L. N. Labzowsky, G. Plunien, T. Beier, and G. Soff, Phys. Rev. A **54**, 2805 (1996).
- [10] V. A. Yerokhin, A. N. Artemyev, V. M. Shabaev, M. M. Sysak, O. M. Zharebtsov, and G. Soff, Phys. Rev. A **64**, 032109 (2001).
- [11] O. M. Zharebtsov, V. M. Shabaev, and V. A. Yerokhin, Phys. Lett. A **277**, 227 (2000).
- [12] A. Schäfer, G. Soff, P. Indelicato, B. Müller, and W. Greiner, Phys. Lett. A **40**, 7362 (1989).
- [13] V. V. Karasiev, L. N. Labzowsky, and A. V. Nefiodov, Phys. Lett. A **172**, 62 (1992).
- [14] R. W. Dunford, Phys. Rev. A **54**, 3820 (1996).
- [15] L. N. Labzowsky, A. V. Nefiodov, G. Plunien, G. Soff, R. Marrus, and D. Liesen, Phys. Rev. A **63**, 054105 (2001).
- [16] M. Gell-Mann and F. Low, Phys. Rev. **84**, 350 (1951).
- [17] J. Sucher, Phys. Rev. **107**, 1448 (1957).
- [18] L. Labzowsky, V. Karasiev, I. Lindgren, H. Persson, and S. Salomonson, Phys. Scr. **T46**, 150 (1993).
- [19] W. H. Furry, Phys. Rev. **81**, 115 (1951).
- [20] L. Labzowsky, G. Klimchitskaya, and Yu. Dmitriev, *Relativistic Effects in the Spectra of Atomic Systems* (Institute of Physics, Bristol, 1993).
- [21] L. N. Labzowsky, J. Phys. B **26**, 1039 (1993).
- [22] M. G. Veselov and L. N. Labzowsky, *Teoriya atoma. Strojenie elektronnykh obolochek* [Theory of atoms. The structure of the electron shells.] (in Russian) (Nauka, Moscow, 1986).
- [23] T. Beier, Phys. Rep. **62**, 052501 (2001).
- [24] W. R. Johnson and G. Soff, At. Data and Nucl. Data Tables **33**, 405 (1985).
- [25] P. Indelicato and P. J. Mohr, Phys. Rev. A **63**, 052507 (2001).
- [26] B. Åsén, S. Salomonson, and I. Lindgren, Phys. Rev. A **65**, 032516 (2002).
- [27] P. J. Mohr, Phys. Rev. A **46**, 4421 (1992).
- [28] P. J. Mohr and G. Soff, Phys. Rev. Lett. **70**, 158 (1993).
- [29] G. Soff and P. J. Mohr, Phys. Rev. A **38**, 5066 (1988).
- [30] T. Beier, A. N. Artemyev, G. Plunien, V. M. Shabaev, G. Soff, and V. A. Yerokhin, Hyperfine Int. **132**, 369 (2001).
- [31] A. N. Artemyev, V. M. Shabaev, and V. A. Yerokhin, Phys. Rev. A **52**, 1884 (1995).
- [32] V. A. Yerokhin, A. N. Artemyev, T. Beier, G. Plunien, V. M. Shabaev, and G. Soff, Phys. Rev. A **60**, 3522 (1999).
- [33] A. N. Artemyev, T. Beier, G. Plunien, V. M. Shabaev, G. Soff, and V. A. Yerokhin, Phys. Rev. A **60**, 45 (1999).
- [34] G. Plunien and G. Soff, Phys. Rev. A **51**, 1119 (1995); **53**, 4614 (1996).
- [35] A. V. Nefiodov, L. N. Labzowsky, G. Plunien, and G. Soff, Phys. Lett. A **222**, 227 (1996).

TABLE I. The values of nuclear root-mean-square radii employed in this work.

$\langle r^2 \rangle^{1/2}$ (fm)	$^{142}_{60}\text{Nd}$	$^{152}_{62}\text{Sm}$	$^{153}_{63}\text{Eu}$	$^{158}_{64}\text{Gd}$	$^{159}_{65}\text{Tb}$	$^{164}_{66}\text{Dy}$	$^{166}_{68}\text{Er}$	$^{174}_{70}\text{Yb}$	$^{202}_{80}\text{Hg}$	$^{231}_{91}\text{Pa}$	$^{238}_{92}\text{U}$	$^{237}_{93}\text{Np}$
	4.914	5.032	5.041	5.090	5.099	5.224	5.165	5.317	5.467	5.700	5.860	5.744

TABLE II. Different contributions to the second-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb								
$\Delta E^{\text{box,irr}}$	-3.929	-3.996	-4.032	-4.069	-4.107	-4.146	-4.230	-4.314
$\Delta E^{\text{cross,irr}}$	0.019	0.021	0.022	0.023	0.024	0.025	0.028	0.030
ΔE^{total}	-3.909	-3.975	-4.010	-4.046	-4.083	-4.120	-4.203	-4.284
Coulomb-Breit								
$\Delta E^{\text{box,irr}}$	-1.195	-1.287	-1.335	-1.384	-1.435	-1.487	-1.595	-1.709
$\Delta E^{\text{box,red}}$	0.492	0.534	0.555	0.577	0.600	0.624	0.673	0.726
$\Delta E^{\text{cross,irr}}$	-0.064	-0.066	-0.066	-0.067	-0.068	-0.071	-0.075	-0.079
$\Delta E^{\text{cross,red}}$	-0.055	-0.062	-0.065	-0.069	-0.072	-0.076	-0.084	-0.093
ΔE^{total}	-0.822	-0.881	-0.911	-0.943	-0.976	-1.010	-1.080	-1.155
Breit-Breit								
$\Delta E^{\text{box,irr}}$	-0.105	-0.119	-0.126	-0.133	-0.141	-0.148	-0.164	-0.183
$\Delta E^{\text{box,red}}$	0.037	0.042	0.045	0.048	0.052	0.055	0.063	0.071
$\Delta E^{\text{cross,irr}}$	0.017	0.018	0.020	0.022	0.025	0.027	0.030	0.033
$\Delta E^{\text{cross,red}}$	0.001	0.002	0.002	0.002	0.002	0.002	0.003	0.003
ΔE^{total}	-0.050	-0.057	-0.059	-0.061	-0.062	-0.064	-0.068	-0.076
Total								
ΔE	-4.781	-4.913	-4.980	-5.049	-5.120	-5.194	-5.350	-5.515

TABLE III. Different contributions to the second-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb				
$\Delta E^{\text{box,irr}}$	-4.845	-5.676	-5.768	-5.867
$\Delta E^{\text{cross,irr}}$	0.046	0.071	0.074	0.077
ΔE^{total}	-4.799	-5.605	-5.695	-5.790
Coulomb-Breit				
$\Delta E^{\text{box,irr}}$	-2.375	-3.391	-3.499	-3.618
$\Delta E^{\text{box,red}}$	1.037	1.516	1.569	1.624
$\Delta E^{\text{cross,irr}}$	-0.108	-0.138	-0.140	-0.145
$\Delta E^{\text{cross,red}}$	-0.146	-0.232	-0.242	-0.252
ΔE^{total}	-1.592	-2.245	-2.312	-2.391
Breit-Breit				
$\Delta E^{\text{box,irr}}$	-0.302	-0.508	-0.528	-0.554
$\Delta E^{\text{box,red}}$	0.127	0.228	0.239	0.251
$\Delta E^{\text{cross,irr}}$	0.055	0.087	0.093	0.098
$\Delta E^{\text{cross,red}}$	0.007	0.013	0.014	0.014
ΔE^{total}	-0.113	-0.181	-0.185	-0.190
Total				
ΔE	-6.504	-8.032	-8.184	-8.371

TABLE IV. Different contributions to the second-order interelectron interaction for the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb								
$\Delta E^{\text{box,irr}}$	-2.748	-2.813	-2.847	-2.882	-2.918	-2.956	-3.035	-3.118
$\Delta E^{\text{cross,irr}}$	0.006	0.007	0.007	0.007	0.008	0.009	0.010	0.012
ΔE^{total}	-2.743	-2.806	-2.840	-2.875	-2.910	-2.947	-3.024	-3.107
Coulomb-Breit								
$\Delta E^{\text{box,irr}}$	-1.162	-1.259	-1.310	-1.363	-1.417	-1.473	-1.587	-1.708
$\Delta E^{\text{box,red}}$	0.048	0.053	0.056	0.059	0.062	0.065	0.071	0.079
$\Delta E^{\text{cross,irr}}$	-0.024	-0.027	-0.029	-0.031	-0.033	-0.035	-0.037	-0.039
$\Delta E^{\text{cross,red}}$	-0.077	-0.084	-0.088	-0.093	-0.097	-0.102	-0.111	-0.122
ΔE^{total}	-1.214	-1.317	-1.372	-1.428	-1.486	-1.545	-1.664	-1.789
Breit-Breit								
$\Delta E^{\text{box,irr}}$	-0.102	-0.123	-0.134	-0.145	-0.156	-0.167	-0.188	-0.209
$\Delta E^{\text{box,red}}$	0.002	0.003	0.003	0.003	0.004	0.004	0.005	0.005
$\Delta E^{\text{cross,irr}}$	-0.013	-0.014	-0.015	-0.015	-0.016	-0.018	-0.020	-0.022
$\Delta E^{\text{cross,red}}$	0.002	0.002	0.003	0.003	0.003	0.003	0.004	0.004
ΔE^{total}	-0.111	-0.132	-0.143	-0.154	-0.165	-0.177	-0.199	-0.222
Total								
ΔE	-4.068	-4.256	-4.354	-4.456	-4.561	-4.670	-4.888	-5.117

TABLE V. Different contributions to the second-order interelectron interaction for the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb				
$\Delta E^{\text{box,irr}}$	-3.636	-4.472	-4.567	-4.667
$\Delta E^{\text{cross,irr}}$	0.022	0.042	0.044	0.047
ΔE^{total}	-3.614	-4.430	-4.523	-4.620
Coulomb-Breit				
$\Delta E^{\text{box,irr}}$	-2.475	-3.670	-3.810	-3.949
$\Delta E^{\text{box,red}}$	0.124	0.199	0.208	0.216
$\Delta E^{\text{cross,irr}}$	-0.053	-0.070	-0.072	-0.073
$\Delta E^{\text{cross,red}}$	-0.187	-0.302	-0.310	-0.323
ΔE^{total}	-2.590	-3.842	-3.985	-4.128
Breit-Breit				
$\Delta E^{\text{box,irr}}$	-0.377	-0.695	-0.743	-0.790
$\Delta E^{\text{box,red}}$	0.011	0.023	0.025	0.026
$\Delta E^{\text{cross,irr}}$	-0.036	-0.064	-0.066	-0.068
$\Delta E^{\text{cross,red}}$	0.008	0.017	0.018	0.020
ΔE^{total}	-0.394	-0.718	-0.766	-0.811
Total				
ΔE	-6.598	-8.991	-9.274	-9.560

TABLE VI. Different contributions to the second-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb								
$\Delta E^{\text{box,irr}}$	-1.535	-1.555	-1.565	-1.575	-1.586	-1.597	-1.620	-1.643
$\Delta E^{\text{cross,irr}}$	0.001	0.002	0.002	0.002	0.002	0.002	0.002	0.003
ΔE^{total}	-1.534	-1.553	-1.563	-1.574	-1.584	-1.595	-1.617	-1.641
Coulomb-Breit								
$\Delta E^{\text{box,irr}}$	-0.006	-0.007	-0.007	-0.007	-0.008	-0.008	-0.008	-0.008
$\Delta E^{\text{box,red}}$	-0.002	-0.003	-0.003	-0.003	-0.003	-0.003	-0.003	-0.004
$\Delta E^{\text{cross,irr}}$	-0.005	-0.006	-0.007	-0.007	-0.007	-0.008	-0.008	-0.008
$\Delta E^{\text{cross,red}}$	0.000	0.000	0.000	0.001	0.001	0.001	0.001	0.001
ΔE^{total}	-0.013	-0.015	-0.016	-0.016	-0.018	-0.018	-0.019	-0.019
Breit-Breit								
$\Delta E^{\text{box,irr}}$	0.001	0.001	0.001	0.002	0.003	0.003	0.003	0.004
$\Delta E^{\text{box,red}}$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\Delta E^{\text{cross,irr}}$	0.004	0.005	0.005	0.006	0.006	0.006	0.007	0.008
$\Delta E^{\text{cross,red}}$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
ΔE^{total}	0.005	0.006	0.007	0.008	0.008	0.009	0.010	0.012
Total								
ΔE	-1.542	-1.562	-1.572	-1.582	-1.593	-1.605	-1.626	-1.648

TABLE VII. Different contributions to the second-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb				
$\Delta E^{\text{box,irr}}$	-1.785	-1.995	-2.017	-2.041
$\Delta E^{\text{cross,irr}}$	0.005	0.008	0.009	0.009
ΔE^{total}	-1.780	-1.986	-2.009	-2.032
Coulomb-Breit				
$\Delta E^{\text{box,irr}}$	-0.010	-0.011	-0.012	-0.012
$\Delta E^{\text{box,red}}$	-0.006	-0.010	-0.011	-0.011
$\Delta E^{\text{cross,irr}}$	-0.009	-0.015	-0.015	-0.015
$\Delta E^{\text{cross,red}}$	0.002	0.004	0.004	0.004
ΔE^{total}	-0.024	-0.032	-0.033	-0.033
Breit-Breit				
$\Delta E^{\text{box,irr}}$	0.004	0.005	0.005	0.006
$\Delta E^{\text{box,red}}$	0.000	0.000	0.000	0.000
$\Delta E^{\text{cross,irr}}$	0.011	0.017	0.017	0.017
$\Delta E^{\text{cross,red}}$	0.001	0.002	0.002	0.002
ΔE^{total}	0.015	0.024	0.024	0.025
Total				
ΔE	-1.789	-1.995	-2.018	-2.040

TABLE VIII. Different contributions to the second-order interelectron interaction for the three-electron configuration $(1s)^2 2s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb								
$\Delta E^{\text{box,irr}}$	-4.267	-4.330	-4.363	-4.397	-4.432	-4.468	-4.546	-4.622
$\Delta E^{\text{cross,irr}}$	0.012	0.013	0.013	0.014	0.015	0.016	0.017	0.019
$\Delta E^{\text{step,irr}}$	-4.046	-4.104	-4.134	-4.165	-4.197	-4.229	-4.298	-4.370
ΔE^{total}	-8.302	-8.421	-8.484	-8.548	-8.614	-8.682	-8.826	-8.974
Coulomb-Breit								
$\Delta E^{\text{box,irr}}$	-0.606	-0.653	-0.678	-0.703	-0.729	-0.755	-0.809	-0.866
$\Delta E^{\text{box,red}}$	0.243	0.263	0.274	0.285	0.296	0.308	0.332	0.357
$\Delta E^{\text{cross,irr}}$	-0.038	-0.041	-0.042	-0.044	-0.045	-0.046	-0.049	-0.051
$\Delta E^{\text{cross,red}}$	-0.027	-0.030	-0.032	-0.033	-0.035	-0.037	-0.041	-0.045
$\Delta E^{\text{step,irr}}$	-0.171	-0.184	-0.191	-0.197	-0.204	-0.212	-0.226	-0.242
$\Delta E^{\text{step,red}}$	-0.041	-0.045	-0.047	-0.049	-0.051	-0.053	-0.058	-0.063
ΔE^{total}	-0.640	-0.690	-0.715	-0.742	-0.768	-0.795	-0.851	-0.909
Breit-Breit								
$\Delta E^{\text{box,irr}}$	-0.051	-0.057	-0.061	-0.064	-0.067	-0.071	-0.078	-0.088
$\Delta E^{\text{box,red}}$	0.018	0.021	0.023	0.025	0.026	0.028	0.031	0.035
$\Delta E^{\text{cross,irr}}$	0.015	0.016	0.017	0.018	0.020	0.022	0.024	0.027
$\Delta E^{\text{cross,red}}$	0.001	0.001	0.001	0.001	0.001	0.001	0.002	0.002
$\Delta E^{\text{step,irr}}$	0.008	0.009	0.009	0.010	0.011	0.012	0.013	0.015
$\Delta E^{\text{step,red}}$	-0.003	-0.003	-0.003	-0.004	-0.004	-0.004	-0.005	-0.006
ΔE^{total}	-0.012	-0.013	-0.013	-0.013	-0.013	-0.013	-0.013	-0.014
Total								
ΔE	-8.954	-9.124	-9.213	-9.303	-9.395	-9.490	-9.690	-9.898

TABLE IX. Different contributions to the second-order interelectron interaction for the three-electron configuration $(1s)^2 2s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb				
$\Delta E^{\text{box,irr}}$	-5.100	-5.830	-5.910	-5.995
$\Delta E^{\text{cross,irr}}$	0.030	0.048	0.050	0.052
$\Delta E^{\text{step,irr}}$	-4.802	-5.455	-5.526	-5.601
ΔE^{total}	-9.872	-11.238	-11.386	-11.543
Coulomb-Breit				
$\Delta E^{\text{box,irr}}$	-1.202	-1.712	-1.771	-1.825
$\Delta E^{\text{box,red}}$	0.509	0.743	0.768	0.795
$\Delta E^{\text{cross,irr}}$	-0.069	-0.092	-0.092	-0.095
$\Delta E^{\text{cross,red}}$	-0.071	-0.111	-0.115	-0.120
$\Delta E^{\text{step,irr}}$	-0.330	-0.453	-0.465	-0.478
$\Delta E^{\text{step,red}}$	-0.094	-0.144	-0.149	-0.155
ΔE^{total}	-1.256	-1.768	-1.824	-1.879
Breit-Breit				
$\Delta E^{\text{box,irr}}$	-0.143	-0.245	-0.253	-0.267
$\Delta E^{\text{box,red}}$	0.064	0.114	0.120	0.126
$\Delta E^{\text{cross,irr}}$	0.044	0.071	0.075	0.078
$\Delta E^{\text{cross,red}}$	0.004	0.009	0.009	0.010
$\Delta E^{\text{step,irr}}$	0.027	0.049	0.052	0.054
$\Delta E^{\text{step,red}}$	-0.010	-0.019	-0.020	-0.021
ΔE^{total}	-0.015	-0.021	-0.017	-0.019
Total				
ΔE	-11.143	-13.028	-13.228	-13.441

TABLE X. Different contributions to the second-order interelectron interaction for the three-electron configuration $(1s)^2 2p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb								
$\Delta E^{\text{box,irr}}$	-8.253	-8.276	-8.294	-8.317	-8.345	-8.378	-8.456	-8.552
$\Delta E^{\text{cross,irr}}$	0.013	0.015	0.016	0.017	0.018	0.019	0.021	0.024
$\Delta E^{\text{step,irr}}$	-5.202	-5.462	-5.591	-5.721	-5.851	-5.981	-6.244	-6.512
ΔE^{total}	-13.443	-13.723	-13.870	-14.022	-14.178	-14.340	-14.679	-15.040
Coulomb-Breit								
$\Delta E^{\text{box,irr}}$	0.226	0.170	0.137	0.102	0.066	0.029	-0.047	-0.131
$\Delta E^{\text{box,red}}$	0.023	0.024	0.026	0.027	0.028	0.029	0.032	0.034
$\Delta E^{\text{cross,irr}}$	-0.026	-0.029	-0.031	-0.033	-0.035	-0.037	-0.040	-0.042
$\Delta E^{\text{cross,red}}$	-0.048	-0.052	-0.054	-0.057	-0.059	-0.061	-0.066	-0.072
$\Delta E^{\text{step,irr}}$	-1.766	-1.823	-1.853	-1.883	-1.915	-1.948	-2.016	-2.088
$\Delta E^{\text{step,red}}$	-0.075	-0.082	-0.086	-0.089	-0.093	-0.097	-0.105	-0.114
ΔE^{total}	-1.666	-1.792	-1.862	-1.934	-2.008	-2.085	-2.242	-2.413
Breit-Breit								
$\Delta E^{\text{box,irr}}$	-0.112	-0.130	-0.138	-0.145	-0.152	-0.158	-0.170	-0.184
$\Delta E^{\text{box,red}}$	0.001	0.001	0.002	0.002	0.002	0.002	0.002	0.003
$\Delta E^{\text{cross,irr}}$	0.004	0.007	0.009	0.011	0.013	0.016	0.023	0.030
$\Delta E^{\text{cross,red}}$	0.002	0.003	0.003	0.003	0.003	0.003	0.004	0.004
$\Delta E^{\text{step,irr}}$	0.119	0.125	0.127	0.130	0.132	0.135	0.140	0.145
$\Delta E^{\text{step,red}}$	0.003	0.003	0.003	0.003	0.004	0.004	0.004	0.005
ΔE^{total}	0.017	0.009	0.006	0.005	0.004	0.003	0.003	0.003
Total								
ΔE	-15.092	-15.506	-15.726	-15.951	-16.182	-16.422	-16.919	-17.450

TABLE XI. Different contributions to the second-order interelectron interaction for the three-electron configuration $(1s)^2 2p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb				
$\Delta E^{\text{box,irr}}$	-9.313	-10.792	-10.968	-11.155
$\Delta E^{\text{cross,irr}}$	0.043	0.078	0.082	0.087
$\Delta E^{\text{step,irr}}$	-7.984	-10.080	-10.309	-10.548
ΔE^{total}	-17.254	-20.794	-21.195	-21.616
Coulomb-Breit				
$\Delta E^{\text{box,irr}}$	-0.619	-1.366	-1.446	-1.532
$\Delta E^{\text{box,red}}$	0.051	0.078	0.081	0.084
$\Delta E^{\text{cross,irr}}$	-0.060	-0.082	-0.078	-0.084
$\Delta E^{\text{cross,red}}$	-0.105	-0.166	-0.166	-0.174
$\Delta E^{\text{step,irr}}$	-2.524	-3.195	-3.269	-3.346
$\Delta E^{\text{step,red}}$	-0.173	-0.278	-0.291	-0.304
ΔE^{total}	-3.430	-5.009	-5.170	-5.357
Breit-Breit				
$\Delta E^{\text{box,irr}}$	-0.389	-0.567	-0.585	-0.622
$\Delta E^{\text{box,red}}$	0.005	0.011	0.012	0.013
$\Delta E^{\text{cross,irr}}$	0.056	0.113	0.118	0.123
$\Delta E^{\text{cross,red}}$	0.009	0.017	0.018	0.019
$\Delta E^{\text{step,irr}}$	0.168	0.189	0.191	0.193
$\Delta E^{\text{step,red}}$	0.009	0.014	0.014	0.015
ΔE^{total}	-0.143	-0.223	-0.232	-0.259
Total				
ΔE	-20.827	-26.027	-26.597	-27.232

TABLE XII. Different contributions to the third-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb-Coulomb	0.010	0.010	0.010	0.010	0.010	0.010	0.010	.010
Coulomb-Coulomb-Breit	0.011	0.011	0.011	0.012	0.012	0.012	0.013	.013
Coulomb-Breit-Breit	0.005	0.005	0.005	0.005	0.005	0.006	0.006	.006
Total	0.026	0.026	0.026	0.027	0.027	0.028	0.029	.029

TABLE XIII. Different contributions to the third-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb-Coulomb	0.012	0.016	0.016	0.017
Coulomb-Coulomb-Breit	0.016	0.022	0.022	0.023
Coulomb-Breit-Breit	0.008	0.011	0.012	0.012
Total	0.036	0.049	0.050	0.052

TABLE XIV. Different contributions to the third-order interelectron interaction for the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb-Coulomb	-0.005	-0.004	-0.004	-0.004	-0.004	-0.003	-0.003	-0.003
Coulomb-Coulomb-Breit	0.011	0.012	0.012	0.013	0.014	0.014	0.015	0.015
Coulomb-Breit-Breit	0.006	0.005	0.005	0.006	0.006	0.006	0.007	0.008
Total	0.013	0.013	0.014	0.015	0.016	0.017	0.019	0.020

TABLE XV. Different contributions to the third-order interelectron interaction for the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb-Coulomb	0.000	0.004	0.004	0.005
Coulomb-Coulomb-Breit	0.021	0.032	0.033	0.034
Coulomb-Breit-Breit	0.013	0.020	0.021	0.022
Total	0.034	0.056	0.058	0.061

TABLE XVI. Different contributions to the third-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb-Coulomb	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001	-0.001
Coulomb-Coulomb-Breit	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
Coulomb-Breit-Breit	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	-0.001	-0.001	-0.001	-0.001	0.000	0.000	0.000	0.000

TABLE XVII. Different contributions to the third-order interelectron interaction for the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb-Coulomb	-0.001	0.000	0.000	0.000
Coulomb-Coulomb-Breit	0.001	0.001	0.001	0.001
Coulomb-Breit-Breit	0.000	0.000	0.000	0.000
Total	0.000	0.001	0.001	0.001

TABLE XVIII. Different contributions to the third-order interelectron interaction for the three-electron configuration $(1s)^22s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb-Coulomb	0.003	0.004	0.004	0.005	0.006	0.006	0.007	0.008
Coulomb-Coulomb-Breit	0.014	0.014	0.015	0.016	0.016	0.016	0.016	0.017
Coulomb-Breit-Breit	0.002	0.003	0.004	0.004	0.005	0.006	0.006	0.006
Total	0.018	0.021	0.023	0.025	0.027	0.028	0.029	0.031

TABLE XIX. Different contributions to the third-order interelectron interaction for the three-electron configuration $(1s)^22s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb-Coulomb	0.015	0.025	0.026	0.027
Coulomb-Coulomb-Breit	0.021	0.028	0.029	0.030
Coulomb-Breit-Breit	0.007	0.011	0.011	0.012
Total	0.043	0.064	0.066	0.069

TABLE XX. Different contributions to the third-order interelectron interaction for the three-electron configuration $(1s)^22p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	65	66	68	70
Coulomb-Coulomb-Coulomb	0.002	0.005	0.007	0.009	0.011	0.013	0.016	0.019
Coulomb-Coulomb-Breit	0.049	0.051	0.053	0.054	0.055	0.056	0.058	0.059
Coulomb-Breit-Breit	0.010	0.011	0.011	0.011	0.011	0.012	0.012	0.012
Total	0.061	0.067	0.071	0.074	0.077	0.080	0.085	0.090

TABLE XXI. Different contributions to the third-order interelectron interaction for the three-electron configuration $(1s)^22p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93
Coulomb-Coulomb-Coulomb	0.041	0.079	0.083	0.088
Coulomb-Coulomb-Breit	0.084	0.116	0.119	0.124
Coulomb-Breit-Breit	0.017	0.030	0.031	0.033
Total	0.142	0.225	0.233	0.245

TABLE XXII. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	Ref.
Zero-order	-13063.004	-14015.106	-14506.589	-15008.567	TW
Nuclear size (NS)	0.928	1.201	1.342	1.519	TW
First-order interelectron interaction	438.812	458.300	468.260	478.371	TW
Second-order interelectron interaction	-4.781 -4.7721	-4.913	-4.980	-5.049	TW [26]
Third-order interelectron interaction	0.026	0.026	0.026	0.027	TW
SE with NS	11.409	12.896	13.700	14.544	[27,28]
VP with NS	-1.621	-1.893	-2.042	-2.201	[24,29]
SE screening	-0.257	-0.289	-0.304	-0.320	[25]
VP screening	0.100	0.113	0.120	0.126	[30]
Recoil	0.055	0.057	0.058	0.059	[31]
Total	-12618.333	-13549.608	-14030.409	-14521.491	TW

TABLE XXIII. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 65$	66	68	70	Ref.
Zero-order	-15521.201	-16044.661	-17124.764	-18250.361	TW
Nuclear size (NS)	1.696	1.968	2.386	3.099	TW
First-order interelectron interaction	488.637	499.061	520.422	542.484	TW
Second-order interelectron interaction	-5.120	-5.194	-5.350	-5.515	TW
Third-order interelectron interaction	0.027	0.028	0.029	0.029	TW
SE with NS	15.427	16.358	18.363	20.589	[27,28]
VP with NS	-2.374	-2.557	-2.962	-3.419	[24,29]
SE screening	-0.335	-0.351	-0.382	-0.413	[25]
VP screening	0.133	0.139	0.164	0.188	[30]
Recoil	0.060	0.061	0.064	0.067	[31]
Total	-15023.050	-15535.148	-16592.030	-17693.252	TW

TABLE XXIV. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2s_{1/2}^1S_0$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93	Ref.
Zero-order	-24622.160	-33320.132	-34215.481	-35130.460	TW
Nuclear size (NS)	9.34	32.50	37.76	40.94	TW
First-order interelectron interaction	665.38	832.758	850.116	868.029	TW
Second-order interelectron interaction	-6.504	-8.032	-8.184 -8.2131	-8.371	TW [26]
Third-order interelectron interaction	0.038	0.049	0.052	0.052	TW
SE with NS	35.3911	62.226	65.4183	68.739	[27,28]
VP with NS	-6.900	-14.632	-15.658	-16.770	[24,29]
SE screening	-0.651	-1.087	-1.127	-1.167	[25]
VP screening	0.354	0.729	0.777	0.825	[30]
Recoil	0.086	0.122	0.127	0.132	[31]
Total	-23925.626	-32415.499	-33286.200	-34178.051	TW

TABLE XXV. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	Ref.
Zero-order	-13063.004	-14015.106	-14506.589	-15008.567	TW
Nuclear size (NS)	0.038	0.052	0.061	0.071	TW
First-order interelectron interaction	447.329	468.642	479.586	490.731	TW
Second-order interelectron interaction	-4.068 -4.0645	-4.256	-4.354	-4.456	TW [3]
Third-order interelectron interaction	0.013	0.013	0.014	0.015	TW
SE with NS	0.298	0.406	0.469	0.539	[27,28]
VP with NS	-0.098	-0.124	-0.139	-0.155	[24,29]
SE screening	-0.112	-0.133	-0.144	-0.155	[25]
VP screening	0.035	0.039	0.042	0.044	[30]
Recoil	0.050	0.051	0.052	0.052	[31]
Total	-12619.519	-13550.416	-14031.002	-14521.881	TW

TABLE XXVI. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 65$	66	68	70	Ref.
Zero-order	-15521.201	-16044.661	-17124.764	-18250.361	TW
Nuclear size (NS)	0.083	0.099	0.129	0.180	TW
First-order interelectron interaction	502.083	513.649	537.458	562.218	TW
Second-order interelectron interaction	-4.561	-4.670	-4.888	-5.117 -5.1140	TW [3]
Third-order interelectron interaction	0.016	0.017	0.019	0.020	TW
SE with NS	0.615	0.700	0.899	1.137	[27,28]
VP with NS	-0.173	-0.193	-0.239	-0.297	[24,29]
SE screening	-0.165	-0.176	-0.197	-0.218	[25]
VP screening	0.047	0.049	0.060	0.070	[30]
Recoil	0.053	0.054	0.055	0.057	[31]
Total	-15023.203	-15535.132	-16591.468	-17692.311	TW

TABLE XXVII. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2p_{1/2}^3P_0$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93	Ref.
Zero-order	-24622.160	-33320.132	-34215.481	-35130.460	TW
Nuclear size (NS)	0.75	3.68	4.41	4.93	TW
First-order interelectron interaction	702.993	902.031	923.176	944.984	TW
Second-order interelectron interaction	-6.598 -6.5959	-8.991	-9.274 -9.2760	-9.560	TW [3]
Third-order interelectron interaction	0.034	0.056	0.058	0.061	TW
SE with NS	3.234	8.772	9.550	10.376	[27,28]
VP with NS	-0.834	-2.451	-2.704	-2.982	[24,29]
SE screening	-0.416	-0.857	-0.897	-0.937	[25]
VP screening	0.144	0.338	0.365	0.391	[30]
Recoil	0.068	0.084	0.086	0.088	[31]
Total	-23922.785	-32417.470	-33290.711	-34183.109	TW

TABLE XXVIII. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	Ref.
Zero-order	-13063.004	-14015.106	-14506.589	-15008.567	TW
Nuclear size (NS)	0.928	1.201	1.342	1.519	TW
First-order interelectron interaction	333.849	347.123	353.854	360.649	TW
Second-order interelectron interaction	-1.542 -1.5455	-1.562	-1.572	-1.582	TW [3]
Third-order interelectron interaction	-1.5459 -0.001	-0.001	-0.001	-0.001	[26] TW
SE with NS	11.409	12.896	13.700	14.544	[27,28]
VP with NS	-1.621	-1.893	-2.042	-2.201	[24,29]
SE screening	-0.196	-0.218	-0.229	-0.240	[25]
VP screening	0.052	0.058	0.061	0.064	[30]
Recoil	0.055	0.057	0.058	0.059	[31]
Total	-12710.071	-13657.445	-14141.418	-14635.756	TW

TABLE XXIX. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 65$	66	68	70	Ref.
Zero-order	-15521.201	-16044.661	-17124.764	-18250.361	TW
Nuclear size (NS)	1.696	1.968	2.386	3.099	TW
First-order interelectron interaction	367.512	374.442	388.524	402.904	TW
Second-order interelectron interaction	-1.593	-1.605	-1.626	-1.648 -1.6548	TW [3]
Third-order interelectron interaction	0.000	0.000	0.000	0.000	TW
SE with NS	15.427	16.358	18.363	20.589	[27,28]
VP with NS	-2.374	-2.557	-2.962	-3.419	[24,29]
SE screening	-0.251	-0.263	-0.285	-0.307	[25]
VP screening	0.067	0.070	0.081	0.092	[30]
Recoil	0.060	0.061	0.064	0.067	[31]
Total	-15140.657	-15656.187	-16720.219	-17828.984	TW

TABLE XXX. Different contributions to the total energy of the two-electron configuration $1s_{1/2}2s_{1/2}^3S_1$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93	Ref.
Zero-order	-24622.160	-33320.132	-34215.481	-35130.460	TW
Nuclear size (NS)	9.34	32.50	37.76	40.94	TW
First-order interelectron interaction	480.13	578.374	588.169	598.188	TW
Second-order interelectron interaction	-1.789 -1.7956	-1.995	-2.018 -2.0203	-2.040	TW [3]
Third-order interelectron interaction	0.000	0.000	-2.0220 0.000	0.000	[26] TW
SE with NS	35.391	62.226	65.418	68.739	[27,28]
VP with NS	-6.914	-14.632	-15.658	-16.770	[24,29]
SE screening	-0.470	-0.754	-0.780	-0.806	[25]
VP screening	0.162	0.310	0.329	0.347	[30]
Recoil	0.086	0.122	0.127	0.132	[31]
Total	-24106.224	-32663.981	-33542.134	-34441.730	TW

TABLE XXXI. Different contributions to the total energy of the three-electron configuration $(1s)^22s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	Ref.
Zero-order	-13063.004	-14015.106	-14506.589	-15008.567	TW
Nuclear size (NS)	0.928	1.201	1.342	1.519	TW
First-order interelectron interaction	720.180	749.835	764.911	780.160	TW
Second-order interelectron interaction	-8.954 -8.953	-9.124	-9.213	-9.303	TW [10]
Third-order interelectron interaction	0.018 0.026	0.021	0.023	0.025	TW [11]
SE with NS	11.409	12.896	13.700	14.544	[27,28]
VP with NS	-1.621	-1.893	-2.042	-2.201	[24,29]
SE screening	-0.820	-0.911	-0.957	-1.002	[32]
VP screening	0.119	0.138	0.147	0.156	[33]
Recoil	0.055	0.057	0.058	0.059	[31]
Total	-12341.690	-13262.886	-13738.620	-14224.610	TW

TABLE XXXII. Different contributions to the total energy of the three-electron configuration $(1s)^2 2s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 65$	66	68	70	Ref.
Zero-order	-15521.201	-16044.661	-17124.764	-18250.361	TW
Nuclear size (NS)	1.696	1.968	2.386	3.099	TW
First-order interelectron interaction	795.587	811.194	842.996	875.598	TW
Second-order interelectron interaction	-9.395	-9.490	-9.690	-9.898 -9.899	TW [10]
Third-order interelectron interaction	0.027	0.028	0.029	0.031 0.039	TW [11]
SE with NS	15.427	16.358	18.363	20.589	[27,28]
VP with NS	-2.374	-2.557	-2.962	-3.419	[24,29]
SE screening	-1.048	-1.093	-1.205	-1.316	[32]
VP screening	0.166	0.175	0.200	0.225	[33]
Recoil	0.060	0.061	0.064	0.067	[31]
Total	-14721.055	-15228.017	-16274.583	-17365.385	TW

TABLE XXXIII. Different contributions to the total energy of the three-electron configuration $(1s)^2 2s_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2s_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93	Ref.
Zero-order	-24622.160	-33320.132	-34215.481	-35130.460	TW
Nuclear size (NS)	9.34	32.50	37.76	40.94	TW
First-order interelectron interaction	795.587	811.194	842.996	875.598	TW
Second-order interelectron interaction	-11.143 -11.147	-13.028	-13.228 -13.226	-13.441	TW [10]
Third-order interelectron interaction	0.043 0.055	0.064	0.066 0.078	0.069	TW [11]
SE with NS	35.391	62.226	65.418	68.739	[27,28]
VP with NS	-6.914	-14.632	-15.658	-16.770	[24,29]
SE screening	-2.063	-3.354	-3.502	-3.650	[32]
VP screening	0.418	0.830	0.882	0.934	[33]
Recoil	0.086	0.122	0.127	0.132	[31]
Nuclear polarization			-0.0377		[34,35]
Total	-23801.415	-3244.210	-33300.658	-34177.909	TW

TABLE XXXIV. Different contributions to the total energy of the three-electron configuration $(1s)^2 2p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 60$	62	63	64	Ref.
Zero-order	-13063.004	-14015.106	-14506.589	-15008.567	TW
Nuclear size (NS)	0.038	0.052	0.061	0.071	TW
First-order interelectron interaction	875.619	913.798	933.291	953.066	TW
Second-order interelectron interaction	-15.092 -15.158	-15.506	-15.726	-15.951	TW [10]
Third-order interelectron interaction	0.061 0.050	0.067	0.071	0.074	TW [11]
SE with NS	0.298	0.406	0.469	0.539	[27,28]
VP with NS	-0.098	-0.124	-0.139	-0.155	[24,29]
SE screening	-0.262	-0.303	-0.324	-0.345	[32]
VP screening	0.044	0.053	0.057	0.061	[33]
Recoil	0.023	0.024	0.025	0.025	[31]
Total	-12202.373	-13116.639	-13588.804	-14071.182	TW

TABLE XXXV. Different contributions to the total energy of the three-electron configuration $(1s)^2 2p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 65$	66	68	70	Ref.
Zero-order	-15521.201	-16044.661	-17124.764	-18250.361	TW
Nuclear size (NS)	0.083	0.099	0.129	0.180	TW
First-order interelectron interaction	973.133	993.500	1035.185	1078.206	TW
Second-order interelectron interaction	-16.182	-16.422	-16.919	-17.450 -17.546	TW [10]
Third-order interelectron interaction	0.077	0.080	0.085	0.090 0.086	TW [11]
SE with NS	0.615	0.700	0.899	1.137	[27,28]
VP with NS	-0.173	-0.193	-0.239	-0.297	[24,29]
SE screening	-0.365	-0.386	-0.442	-0.498	[32]
VP screening	0.066	0.070	0.083	0.095	[33]
Recoil	0.026	0.027	0.028	0.029	[31]
Total	-14563.921	-15067.186	-16105.955	-17188.869	TW

TABLE XXXVI. Different contributions to the total energy of the three-electron configuration $(1s)^2 2p_{1/2}$ (eV). The numbers in the Table present the ionization energy of the $2p_{1/2}$ electron with the opposite sign.

Contribution	$Z = 80$	91	92	93	Ref.
Zero-order	-24622.160	-33320.132	-34215.481	-35130.460	TW
Nuclear size (NS)	0.75	3.68	4.41	4.93	TW
First-order interelectron interaction	1317.19	1642.274	1676.142	1710.926	TW
Second-order interelectron interaction	-20.827 -20.828	-26.027	-26.597 -26.597	-27.232	TW [10]
Third-order interelectron interaction	0.142 0.131	0.225	0.233 0.209	0.245	TW [11]
SE with NS	3.234	8.772	9.550	10.376	[27,28]
VP with NS	-0.834	-2.451	-2.704	-2.982	[24,29]
SE screening	-0.931	-1.860	-1.977	-2.095	[32]
VP screening	0.203	0.483	0.522	0.560	[33]
Recoil	0.038	0.054	0.056	0.057	[31]
Nuclear polarization			-0.0039		[34,35]
Total	-23323.195	-31694.982	-32555.850	-33435.675	TW

TABLE XXXVII. Different theoretical data for the energy levels of two-electron configurations. The numbers in the Table present the ionization energy of the $2s_{1/2}$ or $2p_{1/2}$ electron with the opposite sign, respectively.

Contribution	$Z = 60$	62	63	64	65	66	68	70
$E(2^1S_0)$								
This work	-12618.333	-13549.608	-14030.409	-14521.491	-15023.050	-15535.148	-16592.030	-17693.252
Drake [1]	-12618.188	-13549.412	-14030.191	-14521.251	-15022.780	-15534.938	-16591.716	-17692.948
Plante et al [2]	-12618.629	-13549.909		-14521.817		-15535.580	-16592.446	-17693.778
$E(2^3P_0)$								
This work	-12619.519	-13550.416	-14031.002	-14521.881	-15023.203	-15535.132	-16591.468	-17692.311
Drake [1]	-12619.024	-13549.822	-14030.359	-14521.184	-15022.452	-15534.324	-16590.550	-17691.276
Plante et al [2]	-12619.639	-13550.526		-14521.987		-15535.239	-16591.592	-17692.458
$E(2^3S_1)$								
This work	-12720.071	-13657.445	-14141.418	-14635.756	-15140.657	-15656.187	-16720.219	-17828.984
Drake [1]	-12720.166	-13657.533	-14141.506	-14635.845	-15140.741	-15656.359	-16720.343	-17829.187
Plante et al [2]	-12720.253	-13657.628		-14635.951		-15656.475	-16720.473	-17829.334
$E(2^1S_0) - E(2^3P_0)$								
This work	1.186	0.808	0.593	0.390	0.153	-0.016	-0.562	-0.941
Drake [1]	0.835	0.411	0.168	-0.067	-0.328	-0.614	-1.166	-1.672
Plante et al [2]	1.010	0.617		0.170		-0.341	-0.855	-1.320
$E(2^1S_0) - E(2^3S_1)$								
This work	101.738	107.837	111.009	114.265	117.607	121.039	128.189	135.732
Drake [1]	101.978	108.121	111.315	114.594	117.961	121.421	128.626	136.240
Plante et al [2]	101.624	107.719		114.134		120.895	128.027	135.556
$E(2^3P_0) - E(2^3S_1)$								
This work	100.552	107.029	110.416	113.875	117.454	121.055	128.751	136.673
Drake [1]	101.143	107.710	111.147	114.661	118.290	122.035	129.793	137.911
Plante et al [2]	100.614	107.102		113.963		121.236	128.882	136.876

TABLE XXXVIII. Different theoretical data for the energy levels of two-electron configurations. The numbers in the Table present the ionization energy of the $2s_{1/2}$ or $2p_{1/2}$ electron with the opposite sign, respectively.

Contribution	$Z = 80$	91	92	93
$E(2^1S_0)$				
This work	-23925.626	-32415.499	-33286.200	-34178.051
Drake [1]	-23924.725	-32413.255	-33284.719	-34175.303
Plante et al [2]	-23926.313	-13549.909	-33288.445	
$E(2^3P_0)$				
This work	-23922.785	-32417.470	-33290.711	-34183.109
Drake [1]	-23920.774	-32413.464	-33286.535	-34178.540
Plante et al [2]	-23922.959	-13550.526	-33291.084	
$E(2^3S_1)$				
This work	-24106.224	-32663.981	-33542.134	-34441.730
Drake [1]	-24106.335	-32664.052	-33543.167	-34441.681
Plante et al [2]	-24106.610		-33543.870	
$E(2^1S_0) - E(2^3P_0)$				
This work	-2.841	1.971	4.511	5.058
Drake [1]	-3.951	0.209	1.816	3.237
Plante et al [2]	-3.354		2.640	
$E(2^1S_0) - E(2^3S_1)$				
This work	180.598	248.482	255.934	263.679
Drake [1]	181.610	250.796	258.448	266.378
Plante et al [2]	180.297		255.425	
$E(2^3P_0) - E(2^3S_1)$				
This work	183.439	246.511	251.423	258.621
Drake [1]	185.561	250.587	256.632	263.141
Plante et al [2]	183.651		252.786	




FIG. 1. Feynman graph, describing the first-order interelectron interaction. The double solid lines correspond to bound electrons in the field of the nucleus, the wavy line corresponds to the sum of the Coulomb and Breit (transverse) photons. If $a' = a$ and $b' = b$ the graph is called “direct”, in case $a' = b$, $b' = a$ we call it “exchange” graph. The latter name should be understood in connection with respect to permutation symmetry.




FIG. 2. Feynman graphs describing the second-order interelectron interaction. The graph a) is called “box” and the graph b) is called “cross”. Notations are the same as in Fig. 1. By n_1, n_2 the summation over intermediate states is indicated.




FIG. 3. The third-order “box” Feynman graph. The notations are the same as in Figs. 1, 2. Here the wavy lines with the cross denote the sum of the Coulomb and unretarded Breit interaction.




FIG. 4. The second-order “step” graph for three-electron ions. The notations are the same as in Figs. 1, 2.




FIG. 5. The third-order "step-box" graphs. The wavy line with the cross denotes the sum of the Coulomb and unretarded Breit interactions. Otherwise, the notations are the same as in Figs. 1, 2, 3.